

*Citation for published version:*

Gasperini, D, King, A, Coles, N, Mahon, M & Webster, R 2020, 'Seeking Heteroatom-Rich Compounds: Synthetic and Mechanistic Studies into Iron Catalyzed Dehydrocoupling of Silanes', *ACS Catalysis*, vol. 10, no. 11, pp. 6102-6112. <https://doi.org/10.1021/acscatal.0c01440>

*DOI:*

[10.1021/acscatal.0c01440](https://doi.org/10.1021/acscatal.0c01440)

*Publication date:*

2020

*Document Version*

Peer reviewed version

[Link to publication](https://doi.org/10.1021/acscatal.0c01440)

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *ACS Catalysis*, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <https://pubs.acs.org/doi/10.1021/acscatal.0c01440>

**University of Bath**

## **Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Seeking Heteroatom-Rich Compounds: Synthetic and Mechanistic Studies into Iron Catalyzed Dehydrocoupling of Silanes

*Danila Gasperini,<sup>†</sup> Andrew K. King,<sup>†</sup> Nathan T. Coles,<sup>†</sup> Mary F. Mahon,<sup>†</sup> Ruth L. Webster<sup>†\*</sup>*

<sup>†</sup>Department of Chemistry, University of Bath, Claverton Down, Bath, United Kingdom, BA2 7AY.

## **Abstract**

A detailed synthetic investigation into the dehydrocoupling of silanes with amines, phosphines and alcohols using an iron pre-catalyst (**1**) is presented. We have furnished over 30 examples of aminosilane synthesis along with kinetic studies using MeBnNH and MePhSiH<sub>2</sub> as coupling partners. The kinetic studies suggest a reversible reaction with silane which generates aminosilane and an Fe-hydride dimer that undergoes rate-limiting protonolysis with amine with N–H bond cleavage in the transition state, consistent with a primary KIE of 2.42(3). The presence of dimers as on-cycle intermediates were analyzed in depth. Beyond this we have explored substrate scope of phosphinosilane formation which shows a preferential heterodehydrocoupling to give the phosphinosilane with primary and secondary silanes. Silylethers can also be prepared and alcohols that contain alkene functionality do not show any tendency to reduce the double bond.

## Keywords

Dehydrocoupling, homogeneous catalysis, reaction mechanisms, iron, silanes

## Introduction

Dehydrocoupling of p-block compounds is an attractive method with which to generate hydrogen gas. There has been a great deal of focus in the literature, using a wide range of metal pre-catalysts, on heterodehydrocoupling of amine-boranes, most notably ammonia-borane ( $\text{H}_3\text{N}\cdot\text{BH}_3$ ) because of the high weight% of  $\text{H}_2$  that can be released from this simple molecule.<sup>1</sup> Although there has been focus on dehydropolymerization of amine- and phosphine-boranes for the preparation of materials with novel properties, there are limited examples of dehydrocoupling and dehydropolymerization of silanes with protic substrates such as amines, phosphines and alcohols.

Much of the seminal research into homogeneous catalysis for amino- and phosphinosilane formation via dehydrocoupling was carried out by Harrod,<sup>2</sup> covering several elegant examples using titanocene catalysts,<sup>3,4</sup> and by Eisen using uranium amides for N–Si bond formation.<sup>5</sup> Other examples of aminosilane formation by dehydrocoupling include work from Hill,<sup>6,7</sup> Sarazin<sup>8-11</sup> and Nembenna<sup>12</sup> on Group II catalysis, Crimmin<sup>13</sup> and Guan using  $\text{Ca}$ <sup>14</sup> and  $\text{Y}$ ,<sup>15</sup> Sadow exploring  $\text{Mg}$ <sup>16</sup> and a range of lanthanides,<sup>17</sup> Tsuchimoto with  $\text{Zn}$ <sup>18</sup> and Wright using  $\text{Al}$ .<sup>19</sup> In general, catalytic aminosilane formation is facile, with most examples in the literature being performed at room temperature in a few hours, with the exception of particularly sterically hindered reagents, which often need heating. In contrast the formation of phosphinosilanes tends to require more forcing conditions and there are far fewer examples in the literature. Recent work from Manners has demonstrated that tris(pentafluorophenyl)borane (BCF) can be used in conjunction with tertiary silanes at 100-130 °C to prepare phosphinosilanes along with elegant examples of cyclic

phosphinosilanes when a primary or secondary silane is employed.<sup>20</sup> Waterman has used  $\text{La}[\text{N}(\text{SiMe}_3)_2]_3$ <sup>21</sup> and a Zr catalyst for heterodehydrocoupling<sup>22</sup> along with silylene elimination from silazanes<sup>23</sup> and Stephan has undertaken P–P bond activation to furnish phosphinosilanes using a Rh catalyst.<sup>20, 24</sup> There are few examples of transition metal catalyzed dehydrocoupling methods for silylether formation, with examples in the literature being limited to late transition metals (likely due to oxophilicity in the early to mid-transition metals). Ito and Sawamura have reported using Cu<sup>25</sup> and Au<sup>26, 27</sup> while Yamada has tested a range of acetylacetonate (acac) salts, with  $\text{Cu}(\text{acac})_2$  giving the highest yields (important to note is the very low reactivity obtained with  $\text{Fe}(\text{acac})_2$ ).<sup>28</sup> Sadow has developed a Zn system<sup>29</sup> and Grubbs has reported an attractive NaOH catalyzed method,<sup>30</sup> but most examples are transition metal-free and follow on from Piers' early report of high levels of reactivity using tertiary silanes and a range of alcohols in the presence of BCF catalyst.<sup>31-33</sup>

We saw iron catalyzed heterodehydrocoupling as a remarkably simple route to prepare highly functionalized silane-protected molecules that have a range of applications. In terms of small molecule chemistry, aminosilanes have been used as masked ammonia reagents in Pd-catalyzed aryl-N bond formation, and undergo facile deprotection to reveal the desilylated aniline product<sup>34</sup> and are postulated intermediates in catalytic Staudinger amidation reactions.<sup>35</sup> Phosphinosilanes have been used in Pd-catalyzed cross-coupling by Stille to prepare tertiary phosphines,<sup>36</sup> by Clarke<sup>37</sup> and Yang and Sun<sup>38, 39</sup> in Ni-catalyzed cross-coupling and for C–CN bond activation. Gudat undertook an organocatalytic approach in cross-coupling to prepare P–C bonds from dihalide reagents and  $\text{TMS-PPh}_2$ .<sup>40</sup> More recently phosphinosilanes have been used by Hirano and Miura for the preparation of 1,2-diphosphines<sup>41, 42</sup> and by Daugulis to functionalize benzyne,


utilizing both Si and P components of the reagent.<sup>43</sup> Oro has also used both components of the phosphinosilane for the Ir-catalyzed functionalization of CO<sub>2</sub>.<sup>44</sup>

Silylethers are ubiquitous in protecting group and traceless directing group<sup>45</sup> strategies in organic synthesis. R<sub>2</sub>HSi-protected alcohols can be easily deprotected using standard acidic work-up procedures as demonstrated by Brunner,<sup>46</sup> Nile<sup>47</sup> and Lavastre and Morken.<sup>48</sup> Hartwig used dehydrocoupling to prepare silylethers using an Ir catalyst and the silylether product was then used as a directing group in C–H borylation,<sup>49</sup> Tamao-Fleming oxidation or Hiyama coupling of benzoxasiloles,<sup>50</sup> which is also formed using an Ir catalyst. Herzon has also recently reported elegant studies into C–H functionalization of (+)-Pleuromutilin<sup>51</sup> where the resulting tertiary siloxane was used to undertake an iridium catalyzed C–H bond silylation and subsequent Tamao-Fleming oxidation of the silylated carbon.

Beyond the heterodehydrocoupling of silanes to prepare silyl-protected organic motifs, polysilazanes are used as coatings and precursors to ceramics<sup>52, 53</sup> while polysiloxanes (or silicones) are widely used materials with applications that include elastomers, lubricants, foams and adhesives.<sup>54, 55</sup>

The chemistry that has already been developed using the iron(II) pre-catalyst **1** demonstrates the versatility of this complex, particularly in main group bond transformations that appear to be redox neutral at iron.<sup>56-62</sup> However, we are continually seeking to understand the fundamental bond making and breaking processes involved in catalysis with this complex in order to try to develop new modes of reactivity. We herein present our synthetic and mechanistic investigations into dehydrocoupling and dehydropolymerization to prepare aminosilanes, phosphinosilanes and


silyl ethers (Scheme 1). Only during the preparation of this manuscript have these classes of transformations been reported using an iron catalyst.<sup>63</sup>



**Scheme 1.** The work presented includes a wide substrate scope with over 50 examples of amino- and phosphinosilanes and silyl ethers being presented, along with mechanistic studies.

## Results and Discussion

### 2.1 Aminosilane substrate scope




**Scheme 2.** Varying the primary amine. <sup>a</sup> mono : di.

A range of different primary amines undergo dehydrocoupling with secondary silanes at room temperature (Scheme 2). The reaction tolerates alkyl and aryl amines and works well with diphenyl- and methylphenyl silanes (**2a** to **2k**). Sterically hindered amines can be used, for example *o*-methoxyanisole gives quantitative yield of **2g** while *tert*-butylamine and tritylamine,

although requiring heating to 80 °C, do give excellent yield of product (**2j** and **2k**). Methylphenylsilane is completely consumed, quantitatively forming the disilylated product of 1,4-xylylenediamine (**2i**), while phenylsilane undergoes double amination with aniline (**2m**) in contrast benzylamine, which gives a 70:30 mixture of dimer to monomer (**2l**). No reaction is observed with tertiary silanes such as triphenylsilane and dimethylphenylsilane in conjunction with aniline or morpholine. Interestingly, substrates that contain a double bond also undergo alkene reduction (**2n** to **2s**), which we presume is a consequence of transfer hydrogenation (TH) analogous to our previous report of TH with amines and boranes.<sup>62</sup> We were also able to furnish oligo(silazane) **2t** as an oil. GPC analysis of this species gives a modest  $M_w$  of 855 g mol<sup>-1</sup>. Unfortunately attempts to increase the molecular weight of the polymer, by using very forcing conditions synonymous with condensation polymerization, or dilute conditions associated with step-growth polymerization, did not increase the chain length. Reducing the catalyst loading to 1 mol% and increasing the reaction temperature to 80 °C only led to a modest increase in chain length.<sup>64</sup>





**Scheme 3.** Varying the secondary amine. <sup>a</sup>10 mol% **1**, 80 °C, 48 h; <sup>b</sup>di (shown) : mono by <sup>29</sup>Si{<sup>1</sup>H} NMR, 50 °C.


*N*-Methylbenzylamine and morpholine undergo dehydrocoupling with diphenyl-, diethyl- and methylphenyl silane equally well under our standard reaction conditions, giving quantitative yield of product within 24 h (**3a** to **3d**, **3f** and **3g**, Scheme 3). To demonstrate scalability 5 mmol of morpholine and methylphenylsilane with 2 mol% **1** went to completion after 48h (73% isolated yield). To demonstrate both robustness and scalability, we carried out a standard reaction of morpholine and methylphenylsilane, and after 24 h further portions of amine and silane were added via syringe. This iterative addition process was repeated until 5 mmol of each reagent was added

over the course of 10 days: the catalyst is clearly still active in solution at this time giving complete conversion to **3g** with only 5 mol% **1** after 10 days. Quantitative yield is obtained when *N*-methylbutylamine is coupled to methylphenylsilane (**3i**). Heterocyclic amines piperidine, indoline and tetrahydroquinoline give quantitative yield of the methylphenylsilyl adducts (**3k** to **3m**). Phenylsilane undergoes double amination with both methylbenzylamine and morpholine (**3n** and **3o**).


## 2.2 Aminosilane mechanistic investigations

Intrigued by whether our catalyst would show any similarities to other detailed amine-silane dehydrocoupling kinetics studies (for example Sarazin and Sadow both report  $v = k [\text{amine}]^0 [\text{silane}] [\text{catalyst}]$  (Ba and Mg respectively<sup>8, 16</sup>) whereas Hill reports  $v = k [\text{amine}] [\text{silane}]^0 [\text{catalyst}]$  with Mg/Ca and  $v = k [\text{amine}] [\text{silane}] [\text{catalyst}]^2$  with Ba<sup>6</sup>), we proceeded to undertake *in situ* NMR monitoring studies. We opted to use the reaction of MeBnNH<sub>2</sub> with MePhSiH<sub>2</sub>. Important to note is that the reaction using 3 mol% **1** is 92% complete after around 5 hours at room temperature. Clearly the reaction is facile and, although all amine-silane dehydrocoupling reactions were allowed to react for 24 h, very high levels of conversion (>90%) can be achieved within a few hours, but extended reaction times are needed to achieve quantitative (>99%) conversion to product in most cases (Schemes 2 and 3). The data obtained suggest that the reaction has a first order dependence on the Fe-center when using **1** as reaction pre-catalyst (Figure 1a). We then studied the transformation by varying the concentration of silane. As can be seen, the reaction is independent of silane concentration (Figure 1b). As expected, with 0.5 equivalents of silane around 50% yield of product is obtained (yellow squares). In contrast, a doubling of amine loading results in an approximate doubling of the reaction rate (Figure 1c) and so a first order dependence in amine is observed.


a)



b)




c)



**Figure 1.** a) Plot showing first order relationship for formation of **3f** at different pre-catalyst loadings (6 mol%, 5 mol%, 4 mol%, 3 mol%). Conditions: 0.5 mmol silane, 0.5 mmol amine, 0.5 mL C<sub>6</sub>D<sub>6</sub>, RT. b) Reaction profiles showing the formation of MePhSi–NMeBn, **3f**, at different silane loadings (○ 2 equiv., ● 1.5 equiv., ■ 1 equiv., □ 0.5 equiv.) Conditions: 3 mol% **1**, 0.5 mmol amine, 0.5 mL C<sub>6</sub>D<sub>6</sub>, RT. c) Plot showing first order relationship for formation of **3f** at different amine loadings (2 equiv., 1.5 equiv., 1 equiv., 0.5 equiv.). Conditions: 3 mol% **1**, 0.5 mmol silane, 0.5 mL C<sub>6</sub>D<sub>6</sub>, RT.

Monitoring the paramagnetic region of a <sup>1</sup>H NMR spectrum for a catalytic reaction shows that iron amido complex **4** is observed throughout. Interestingly, **1** is also observed throughout the reaction. Starting from **1**, using an analogous method to that reported by Holland,<sup>65</sup> **4** was prepared and isolated. Use of **4** in catalysis shows that the lengthy kinetic induction period that is observed with **1** is circumvented and 92% conversion to product is obtained after 4 h, compared to 7 h when using **1** (there is 50% conversion after 90 minutes using **4** and 50% conversion after 150 minutes with **1**). This also suggests that **4** is likely to be an on-cycle species in catalysis. The reaction is first order in **4**, which we attribute to this species only contributing one Fe-center to the key

intermediate (**5/5'** *vide infra*).<sup>64</sup> Holland and co-workers have shown previously that in both the solid state and in solution iron hydride species with a methyl backbone  $\beta$ -diketiminato ancillary ligand exists as the dimeric hydride complex **5**.<sup>66</sup> At 298 K this dimeric species is observed by <sup>1</sup>H NMR in our reaction.<sup>67</sup> Preparation, isolation and kinetics studies with **5** give a first order fit.<sup>64</sup> This was unexpected, but similar results have been reported by Holland when investigating B–C bond cleavage mediated by **5**.<sup>68</sup> This first order fit is attributed to the presence of the open isomer (see **5'**, Scheme 4), which is in rapid equilibrium with **5**. If the amine and the iron hydride dimer (**5**) are involved in the rate-limiting step with *only one Fe–H bond* being broken during this step then only one equivalent of amine should react and therefore we would expect a first order dependence in HNMeBn when **5** is employed directly in catalysis, which is the case.<sup>64</sup> We therefore postulate that the rate limiting transition state involves the formation of a cyclic species, possibly of the form **5'**<sub>TS</sub>, followed by release of hydrogen gas and formation of **4** and **5**. Collapse of **5'**<sub>TS</sub> might proceed via a mixed hydride-amido dimer such as **6**. We have evidence for the ability for such species to form; reacting **5** with H<sub>2</sub>N<sup>n</sup>Bu and MePhSiH<sub>2</sub> **6** is immediately observed by <sup>1</sup>H NMR and X-ray quality crystals were readily obtained (Figure 2).




**Figure 2.** Mixed hydride-amido dimer (**6**) isolated from reaction of  $^n\text{BuNH}_2$ ,  $\text{PhMeSiH}_2$  and **5**.<sup>64</sup> Ellipsoids are represented at 50%. Except for the hydride ligand and N-H, all hydrogen atoms, solvent, disorder and the 2,6-diisopropyl groups have been omitted for clarity.

In order to probe the nuclearity of the species present in solution, DOSY experiments were performed<sup>69</sup> on a mixture of **5** and  $^n\text{BuNH}_2$  in  $\text{C}_6\text{D}_6$ ; disappearance of **5** is observed with  $\text{H}_2$  release and formation of dimers ( $D = 5.04\text{--}5.48 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$ ), primarily consisting of complex **6** but formation of other mixed dimeric species cannot be ruled out ( $r_{\text{solution}} = 8.2 \text{ \AA}$ ).<sup>70</sup> Pleasingly when reacting pre-catalyst **1** with a mixture of amine ( $^n\text{BuNH}_2$ ) and  $\text{PhMeSiH}_2$  in  $\text{C}_6\text{D}_6$  multiple species were observed; the slower peaks were analyzed and showed similar diffusion coefficients ( $D = 4.50\text{--}5.95 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$ ) to the experiment above and a large solution hydrodynamic radius ( $r_{\text{solution}} = 8.5 \text{ \AA}$ ) synonymous with dimers. These results are in line with the appearance of dimer **5** during

kinetic  $^1\text{H}$  NMR analysis and the order in **5** which suggests it exists as an active dimer throughout catalysis. Furthermore, calculation of the hydrodynamic radius of **6** using single crystal data<sup>71</sup> is very similar to the solution data obtained ( $r_{\text{solid}} = 9.2 \text{ \AA}$ )

Combined, these data have allowed us to postulate a preliminary catalytic cycle (Scheme 4). The steps in the catalytic cycle are in-line with reports from Sadow,<sup>16</sup> Sarazin<sup>8</sup> and Hill<sup>6</sup> in that catalyst activation forms an initial metal-amido species which then undergoes metal-hydride formation and release of the aminosilane product. Unlike the report from Sarazin, where hydride transfer is rate limiting, or Sadow where attack of silane by an Mg-amido species is rate limiting, the subsequent proton transfer between *N*-methylbenzylamine and iron-hydride is probably the slow step in our reaction.



**Scheme 4.** Postulated mechanism based on NMR studies.


Deuterium labelling using  $d_2$ -aniline as an easily accessible substrate gives a primary KIE for the formation of **2f** of 2.42(3),<sup>64</sup> which may indicate proton transfer is occurring through a non-linear transition state. This also supports our proposed catalytic cycle where the reaction is first order in amine. We have also undertaken Arrhenius and Eyring analysis of catalysis mediated by **1**:  $E_a = 54.59 \text{ kJ mol}^{-1} \pm 0.03$ ;  $\Delta H^\ddagger = 12.43 \text{ kcal mol}^{-1} \pm 0.02$ ;  $\Delta S^\ddagger = -5.13 \text{ cal mol}^{-1} \text{ K}^{-1} \pm 0.01$ ;



$\Delta G^\ddagger = 13.95 \text{ kcal mol}^{-1} \pm 0.2$ . This is in line with reports using Mg<sup>6</sup> (13.9 kcal mol<sup>-1</sup>) and Ba<sup>8</sup> (15.6 kcal mol<sup>-1</sup>) systems. The latter study from Sarazin also reports similar  $E_a$  and  $\Delta G^\ddagger$  to our system. The importance of  $\Delta H^\ddagger$  to the activation energy barrier for our process is highlighted by the primary KIE value. We can rationalize the low  $\Delta S^\ddagger$  obtained by looking at the process of **5/5'** and thus onward formation of **5'**ts. In these steps an Fe–H bond is broken and H–H bond begins to form. Akin to work from Holland on the likelihood of **5'** to be a reaction intermediate, this ‘interchange mechanism’ is often linked to very low  $\Delta S^\ddagger$ .<sup>67</sup>

On closer inspection of the deuterium labelling study we can report another intriguing result. Reaction of *d*<sub>2</sub>-aniline with methylphenylsilane gives a mixture of PhDN–SiHMePh (**d<sub>1</sub>-2f**) and PhDN–SiDMePh (**d<sub>2</sub>-2f**) in a 77:23 ratio (Scheme 5a). In the absence of catalyst there is no H/D exchange and no dehydrocoupling (to give product(s) of the form **2f**, Scheme 5b). To determine whether the HD released from reaction of *d*<sub>2</sub>-aniline with methylphenylsilane is responsible for silane deuteration, presumably due to HD activation by iron catalyst, we performed the standard reaction under an atmosphere of D<sub>2</sub> (Scheme 5c). Reaction in a J-Young NMR tube gives a small amount of Si–D product (**d<sub>1</sub>-Si-2f**), but no deuterium incorporation at nitrogen takes place (in the aniline starting material or aminosilane product). A standard reaction, of the form presented in Scheme 5a, will generate an estimated 6 bar of H<sub>2</sub>/HD/D<sub>2</sub> pressure in a J-Young NMR tube (based on the Ideal Gas Law), whereas the reactions in Scheme 5c are back-pressurized with 1 bar D<sub>2</sub>, hence the difference in J-Young NMR tube deuterium incorporation results. The same reaction in a Schlenk tube with a large headspace back-filled with 1 atm. D<sub>2</sub> gives 82% **d<sub>1</sub>-Si-2f**. These data indicate that a greater number of equivalents of D<sub>2</sub> leads to more D-incorporation: the same reaction under a D<sub>2</sub> atmosphere was performed in the same scale in a sealed Schlenk (25 mL) without stirring and it showed 87% deuterium incorporation, suggesting that mass transfer is not


influencing the reaction outcome. N–H bond activation is likely more challenging than other steps in the catalytic cycle; in-keeping with our postulated catalytic cycle. **2f** does not show any deuteration when exposed to 1 atm. D<sub>2</sub> (Scheme 5d): the iron catalyst is not able to further functionalize **2f**. Repeating this reaction of **2f** in the presence of one equivalent of *d*<sub>2</sub>-aniline, 1 atm. D<sub>2</sub> and 5 mol% **1** gives a 1:1 mixture of **2f** and *d*<sub>1</sub>-N-**2f**, therefore indicating that N-Si bond formation is reversible (Scheme 5e). Furthermore H/D exchange is not observed when **2f** is mixed with *d*<sub>2</sub>-aniline. Si–H bond deuteration is not observed under these conditions indicating that silazane formation is fast and reversible, precluding any deuteration of free silane released in the reverse reaction. This also suggests that silane deuteration occurs via a different catalytic cycle. Taking **1**, H<sub>2</sub>NPh (to allow catalyst activation) and methylphenylsilane under an atmosphere of D<sub>2</sub> does not show deuterium-incorporation at nitrogen, indicating that direct formation of iron(II)-deuteride from iron(II)-amido does not take place. Reaction of methylphenylsilane, D<sub>2</sub> and **1** (5 mol%) gives 9% MePhSiHD further indicating that H/D exchange via Si–H activation is feasible in the presence of **1** and this process is currently under investigation.



**Scheme 5.** Studies into deuterium incorporation into **2f**.


Based on the combined experimental and literature data an updated catalytic cycle for reaction of  $\text{PhNH}_2$  with  $\text{MePhSiH}_2$  in the presence of  $\text{D}_2$  and incorporating H/D exchange can be proposed

(Scheme 6). The open-isomer was invoked as possible mechanistic scenario for hydride ligand exchange with  $D_2$  from **5** to **5'-D** by Holland and co-workers.<sup>65</sup> The catalytic cycle for Si-D bond formation could then proceed in a similar manner to that of Si-N bond formation, in that there is a fast reversible equilibrium between **5'-D** and the open isomer, reaction with amine releases  $H_2$  or HD and forms **4**, it would be at this point that fast reversible equilibrium between silane and **5'-D** allows deuteration of the silane.



**Scheme 6.** Revised catalyst cycle for dehydrocoupling demonstrating that, starting from a common intermediate (**5'**) a second cycle that could account for Si-D bond formation due to H/D exchange in the presence of  $D_2$ .

### 2.3 Phosphinosilane substrate scope



**Scheme 7.** Scope of phosphinosilane reactivity.


Given the reaction scope achieved with amines, we expected to achieve good reactivity with phosphines. When we began our optimization studies it was immediately clear that phosphine-silane dehydrocoupling is a more challenging target for pre-catalyst **1**: in order to obtain high conversion to the dehydrocoupled product heating to 70 °C is necessary (Scheme 7). For comparison Waterman has heated to 90 °C and Manners to 100 °C with their Zr<sup>22</sup> and BCF<sup>20</sup> catalyzed processes. Also apparent in our catalysis is the potential for competitive phosphine homodehydrocoupling to take place; this is not unexpected as we have previously reported diphosphane synthesis via dehydrocoupling using comparable reaction conditions.<sup>57</sup> Homodehydrocoupling becomes competitive with the desired heterodehydrocoupling reaction when tertiary silanes (triphenyl- and methyldiphenyl-) are employed (**7g** to **7j**) and in the case of diphenylphosphine, complete conversion to tetraphenyldiphosphane (Ph<sub>2</sub>P–PPh<sub>2</sub>) is observed (**7g** and **7i**). When dicyclohexylphosphine is reacted with tertiary silanes, although the transformation is completely selective for homodehydrocoupling, only trace amount of product is observed. This

is in-line with our previous report on phosphine dehydrocoupling i.e. we do not believe that the silane plays an active role in assisting homodehydrocoupling. Moreover, under the catalytic reaction conditions, homodehydrocoupled product  $(\text{Ph}_2\text{P})_2$  does not react with  $\text{Ph}_2\text{SiH}_2$  even after 3 days at 80 °C: our system does not proceed via activation of the P–P bond (as achieved by Manners<sup>20</sup> and Stephan<sup>24</sup>) but rather through activation of the P–H bond. In contrast, when secondary silanes are employed, dicyclohexylphosphine gives much better selectivity for the desired heterodehydrocoupled product. 99:1 Hetero:homodehydrocoupled product is obtained with dicyclohexylphosphine with both diphenylsilane and methylphenyl silane (**7b** and **7d**), whereas this ratio drops to 85:15 and 81:19 respectively when  $\text{HPPH}_2$  is employed (**7a** and **7c**). Again, this is a direct reflection of the ease with which the phosphines homodehydrocouple. Similar to Harrod's report using titanocenes, phenylsilane only reacts once with  $\text{HPPH}_2$  and  $\text{HPCy}_2$  yielding  $\text{R}_2\text{PSiPhH}_2$  selectively.

## 2.4 Silylether substrate scope

The inherent oxophilicity of Fe(II) does not appear to be limiting when alcohols are employed as coupling partners. However, the nature of the alcohol appears to be important and there is an interesting divergence in reactivity whereby phenol only gives a trace amount of product (**8a**, Scheme 8) whereas aliphatic alcohols give almost quantitative yield of siloxane (**8b** to **8g**, with the exception of methanol which only gives 35% **8c**). If the silane is changed from methylphenylsilane to phenylsilane then good levels of mono-silylated phenolic product is achieved (**8m**). Returning to focus on reactivity achieved with methylphenylsilane, 2-phenylglycinol shows interesting selectivity with the alcohol reacting preferentially (**8h**). Elaborate organic motifs can undergo heterodehydrocoupling to generate the silylated product including (–)-menthol (**8g**), cholesterol

(**8i**), linalool (**8j**),  $\beta$ -citronellol (**8k**) with no evidence for double bond functionalization or isomerization. Phenylsilane reacts with benzylalcohol and benzhydrol quantitatively (**8n** and **8o**) and with pinacol alcohol to give the dioxasilolane (**8p**). Oligosiloxane, **8q**, derived from phenylsilane and diethylene glycol was also prepared as a solid with very narrow PDI and *DP* of 4.5. DSC shows two  $T_m$  peaks (245.7 °C and 256.2 °C) which may indicate the presence of both linear and cyclic oligomers.



**Scheme 8.** A range of alcohols can be protected as silylethers using our heterodehydrocoupling methodology. <sup>a</sup>71% achieved after 24 h at 50 °C; <sup>b</sup>intractable mixture of mono- and di-substituted silane and also likely to contain functionalization/isomerization of double bond(s); <sup>c</sup>mono:di; <sup>d</sup>80 °C, 24 h.

### 3. Conclusions

In summary, we have presented a detailed study into the reactivity of a simple iron pre-catalyst **1** towards silane heterodehydrocoupling. Amines react well with secondary silanes, which is complementary to the state-of-the-art in the literature, and give good selectivity for mono-substituted product. A detailed mechanistic study was undertaken which showed that the rate-limiting step was likely to be protonolysis of an iron hydride dimer to release hydrogen and generate an iron-amido. We believe we have provided strong evidence for the presence of iron dimers being active on-cycle species during catalysis. Deuterium labelling studies show a modest primary KIE that support rate limiting protonolysis and suggest that the RLS proceeds via a non-linear transition state. Unexpectedly, Si–D formation in the presence of D<sub>2</sub> appears to be more favorable than N–D formation. However, this is in-line with kinetics results. Finally, we have shown that phosphinosilane formation is possible under mild conditions, while we have also generated a diverse range of structurally complex silylethers: valuable building blocks in organic synthesis.

#### ASSOCIATED CONTENT

##### Supporting Information.

The following files are available free of charge.

Analysis data and NMR spectra for all products (PDF). CCDC 1978051 contains the



supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk/structures>

## AUTHOR INFORMATION

### Corresponding Author

\* R.L.Webster@bath.ac.uk

## ACKNOWLEDGMENT

The EPSRC is thanked for funding and Dr John Lowe is thanked for assistance with DOSY analysis. Many thanks are also extended to Professor Stuart Macgregor and Dr Sam Neale for their efforts with DFT analysis.

## REFERENCES

1. Leita, E. M.; Jurca, T.; Manners, I., Catalysis in service of main group chemistry offers a versatile approach to p-block molecules and materials. *Nat. Chem.* **2013**, 5, 817-829.
2. Gauvin, F.; Harrod, J. F.; Woo, H. G., Catalytic Dehydrocoupling: A General Strategy for the Formation of Element–Element Bonds. In *Adv. Organomet. Chem.*, Stone, F. G. A.; West, R., Eds. Academic Press: **1998**; Vol. 42, pp 363-405.
3. Shu, R. H.; Hao, L. J.; Harrod, J. F.; Woo, H. G.; Samuel, E., Heterodehydrocoupling of phosphines and silanes catalyzed by titanocene: A novel route to the formation of Si-P bonds. *J. Am. Chem. Soc.* **1998**, 120, 12988-12989.
4. Harrod, J. F., Catalysis of reactions of Si-H by titanocene and its derivatives. *Coord. Chem. Rev.* **2000**, 206, 493-531.
5. Wang, J. X.; Dash, A. K.; Berthet, J. C.; Ephritikhine, M.; Eisen, M. S., Dehydrocoupling reactions of amines with silanes catalyzed by [(Et<sub>2</sub>N)<sub>3</sub>U][BPh<sub>4</sub>]. *J. Organomet. Chem.* **2000**, 610, 49-57.
6. Hill, M. S.; Liptrot, D. J.; MacDougall, D. J.; Mahon, M. F.; Robinson, T. P., Heterodehydrocoupling of silanes and amines by heavier alkaline earth catalysis. *Chem. Sci.* **2013**, 4, 4212-4222.
7. Morris, L. J.; Whittell, G. R.; Eloi, J.-C.; Mahon, M. F.; Marken, F.; Manners, I.; Hill, M. S., Ferrocene-Containing Polycarbosilazanes via the Alkaline-Earth-Catalyzed Dehydrocoupling of Silanes and Amines. *Organometallics* **2019**, 38, 3629-3648.

8. Bellini, C.; Carpentier, J.-F.; Tobisch, S.; Sarazin, Y., Barium-Mediated Cross-Dehydrocoupling of Hydrosilanes with Amines: A Theoretical and Experimental Approach. *Angew. Chem. Int. Ed.* **2015**, *54*, 7679-7683.
9. Bellini, C.; Orione, C.; Carpentier, J.-F.; Sarazin, Y., Tailored Cyclic and Linear Polycarbosilazanes by Barium-Catalyzed N–H/H–Si Dehydrocoupling Reactions. *Angew. Chem. Int. Ed.* **2016**, *55*, 3744-3748.
10. Bellini, C.; Dorcet, V.; Carpentier, J.-F.; Tobisch, S.; Sarazin, Y., Alkaline-Earth-Catalysed Cross-Dehydrocoupling of Amines and Hydrosilanes: Reactivity Trends, Scope and Mechanism. *Chem. Eur. J.* **2016**, *22*, 4564-4583.
11. Bellini, C.; Roisnel, T.; Carpentier, J.-F.; Tobisch, S.; Sarazin, Y., Sequential Barium-Catalysed N–H/H–Si Dehydrogenative Cross-Couplings: Cyclodisilazanes versus Linear Oligosilazanes. *Chem. Eur. J.* **2016**, *22*, 15733-15743.
12. Baishya, A.; Peddaraio, T.; Nembenna, S., Organomagnesium amide catalyzed cross-dehydrocoupling of organosilanes with amines. *Dalton Trans.* **2017**, *46*, 5880-5887.
13. Nako, A. E.; Chen, W.; White, A. J. P.; Crimmin, M. R., Yttrium-Catalyzed Amine–Silane Dehydrocoupling: Extended Reaction Scope with a Phosphorus-Based Ligand. *Organometallics* **2015**, *34*, 4369-4375.
14. Li, N.; Guan, B.-T., A Dialkyl Calcium Carbene Adduct: Synthesis, Structure, and Catalytic Cross-Dehydrocoupling of Silanes with Amines. *Eur. J. Inorg. Chem.* **2019**, *2019*, 2231-2235.
15. Li, N.; Guan, B.-T., Yttrium–Benzyl Complexes Bearing Chiral Iminophosphonamide Ligands: Synthesis and Application in Catalytic Asymmetric Amine-Silane Dehydrocoupling Reactions. *Adv. Synth. Catal.* **2017**, *359*, 3526-3531.
16. Dunne, J. F.; Neal, S. R.; Engelkemier, J.; Ellern, A.; Sadow, A. D., Tris(oxazolinyl)boratomagnesium-Catalyzed Cross-Dehydrocoupling of Organosilanes with Amines, Hydrazine, and Ammonia. *J. Am. Chem. Soc.* **2011**, *133*, 16782-16785.
17. Pindwal, A.; Ellern, A.; Sadow, A. D., Homoleptic Divalent Dialkyl Lanthanide-Catalyzed Cross-Dehydrocoupling of Silanes and Amines. *Organometallics* **2016**, *35*, 1674-1683.
18. Yonekura, K.; Iketani, Y.; Sekine, M.; Tani, T.; Matsui, F.; Kamakura, D.; Tsuchimoto, T., Zinc-Catalyzed Dehydrogenative Silylation of Indoles. *Organometallics* **2017**, *36*, 3234-3249.
19. Allen, L. K.; Garcia-Rodriguez, R.; Wright, D. S., Stoichiometric and catalytic Si–N bond formation using the p-block base Al(NMe<sub>2</sub>)<sub>3</sub>. *Dalton Trans.* **2015**, *44*, 12112-12118.
20. Wu, L. P.; Chitnis, S. S.; Jiao, H. J.; Annibale, V. T.; Manners, I., Non-Metal-Catalyzed Heterodehydrocoupling of Phosphines and Hydrosilanes: Mechanistic Studies of B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>-Mediated Formation of P–Si Bonds. *J. Am. Chem. Soc.* **2017**, *139*, 16780-16790.
21. Cibuzar, M. P.; Waterman, R., Si–N Heterodehydrocoupling with a Lanthanide Compound. *Organometallics* **2018**, *37*, 4395-4401.
22. Roering, A. J.; MacMillan, S. N.; Tanski, J. M.; Waterman, R., Zirconium-catalyzed heterodehydrocoupling of primary phosphines with silanes and germanes. *Inorg. Chem.* **2007**, *46*, 6855-6857.
23. Erickson, K. A.; Cibuzar, M. P.; Mucha, N. T.; Waterman, R., Catalytic N–Si coupling as a vehicle for silane dehydrocoupling via  $\alpha$ -silylene elimination. *Dalton Trans.* **2018**, *47*, 2138-2142.

24. Greenberg, S.; Stephan, D. W., Stoichiometric and catalytic activation of P-H and P-P bonds. *Chem. Soc. Rev.* **2008**, *37*, 1482-1489.
25. Ito, H.; Watanabe, A.; Sawamura, M., Versatile Dehydrogenative Alcohol Silylation Catalyzed by Cu(I)-Phosphine Complex. *Org. Lett.* **2005**, *7*, 1869-1871.
26. Ito, H.; Takagi, K.; Miyahara, T.; Sawamura, M., Gold(I)-Phosphine Catalyst for the Highly Chemoselective Dehydrogenative Silylation of Alcohols. *Org. Lett.* **2005**, *7*, 3001-3004.
27. Ito, H.; Saito, T.; Miyahara, T.; Zhong, C.; Sawamura, M., Gold(I) Hydride Intermediate in Catalysis: Dehydrogenative Alcohol Silylation Catalyzed by Gold(I) Complex. *Organometallics* **2009**, *28*, 4829-4840.
28. Yasuhiko, G.; Yoshihiro, Y.; Taketo, I.; Tohru, Y., Convenient and Selective Preparation of Mono-alkoxyphenylsilanes from Phenylsilane and Alcohols. *Chem. Lett.* **2006**, *35*, 714-715.
29. Mukherjee, D.; Thompson, R. R.; Ellern, A.; Sadow, A. D., Coordinatively Saturated Tris(oxazolinyl)borato Zinc Hydride-Catalyzed Cross Dehydrocoupling of Silanes and Alcohols. *ACS Catal.* **2011**, *1*, 698-702.
30. Toutov, A. A.; Betz, K. N.; Haibach, M. C.; Romine, A. M.; Grubbs, R. H., Sodium Hydroxide Catalyzed Dehydrocoupling of Alcohols with Hydrosilanes. *Org. Lett.* **2016**, *18*, 5776-5779.
31. Blackwell, J. M.; Foster, K. L.; Beck, V. H.; Piers, W. E., B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>-Catalyzed Silation of Alcohols: A Mild, General Method for Synthesis of Silyl Ethers. *J. Org. Chem.* **1999**, *64*, 4887-4892.
32. Könczöl, L.; Kawachi, A.; Szieberth, D., Mechanism of Dehydrogenative Condensation of (o-borylphenyl)hydrosilanes with Methanol. *Organometallics* **2012**, *31*, 120-125.
33. Strašák, T.; Sýkora, J.; Lamač, M.; Kubišta, J.; Horáček, M.; Gyepes, R.; Pinkas, J., Reactivity of a Titanocene Pendant Si-H Group toward Alcohols. Unexpected Formation of Siloxanes from the Reaction of Hydrosilanes and Ph<sub>3</sub>COH Catalyzed by B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>. *Organometallics* **2013**, *32*, 4122-4129.
34. Lombardi, C.; Mitchell, D.; Rodriguez, M. J.; Organ, M. G., Pd-PEPPSI-IPentCl-Catalyzed Amination Using Aminotriphenylsilane as an Ammonia Surrogate. *Eur. J. Org. Chem.* **2017**, *2017*, 1510-1513.
35. Andrews, K. G.; Denton, R. M., A more critical role for silicon in the catalytic Staudinger amidation: silanes as non-innocent reductants. *Chem. Commun.* **2017**, *53* (57), 7982-7985.
36. Tunney, S. E.; Stille, J. K., Palladium-catalyzed Coupling of Aryl Halides with (Trimethylstannyl)diphenylphosphine and (Trimethylsilyl)diphenylphosphine. *J. Org. Chem.* **1987**, *52*, 748-753.
37. Clarke, M. L.; Guy Orpen, A.; Pringle, P. G.; Turley, E., Organometallic chemistry of bromodifluoromethyl substituted phosphines. The development of a novel nickel catalysed P-C bond forming reaction. *Dalton Trans.* **2003**, 4393-4394.
38. Sun, M.; Zhang, H.-Y.; Han, Q.; Yang, K.; Yang, S.-D., Nickel-Catalyzed C-P Cross-Coupling by C-CN Bond Cleavage. *Chem. Eur. J.* **2011**, *17*, 9566-9570.
39. Sun, M.; Zang, Y.-S.; Hou, L.-K.; Chen, X.-X.; Sun, W.; Yang, S.-D., Convenient Formation of Triarylphosphines by Nickel-Catalyzed C-P Cross-Coupling with Aryl Chlorides. *Eur. J. Org. Chem.* **2014**, *2014*, 6796-6801.
40. Burck, S.; Förster, D.; Gudat, D., Phosphorus-carbon bond formation catalysed by electrophilic N-heterocyclic phosphines. *Chem. Commun.* **2006**, 2810-2812.

41. Okugawa, Y.; Hirano, K.; Miura, M., Copper-Catalyzed Vicinal Diphosphination of Styrenes: Access to 1,2-Bis(diphenylphosphino)ethane-Type Bidentate Ligands from Olefins. *Angew. Chem. Int. Ed.* **2016**, *55*, 13558-13561.
42. Otomura, N.; Okugawa, Y.; Hirano, K.; Miura, M., vic-Diphosphination of Alkenes with Silylphosphine under Visible-Light-Promoted Photoredox Catalysis. *Org. Lett.* **2017**, *19*, 4802-4805.
43. Mesgar, M.; Nguyen-Le, J.; Daugulis, O., New Hindered Amide Base for Aryne Insertion into Si-P, Si-S, Si-N, and C-C Bonds. *J. Am. Chem. Soc.* **2018**, *140*, 13703-13710.
44. Julián, A.; Polo, V.; Fernández-Alvarez, F. J.; Oro, L. A., Iridium-NSiN catalyzed formation of silylphosphinecarboxylates from the reaction of CO<sub>2</sub> with P(SiMe<sub>3</sub>)R<sub>2</sub> (R = Ph, Cy). *Catal. Sci. Tech.* **2017**, *7*, 1372-1378.
45. Mewald, M.; Schiffner, J. A.; Oestreich, M., A New Direction in C-H Alkenylation: Silanol as a Helping Hand. *Angew. Chem. Int. Ed.* **2012**, *51*, 1763-1765.
46. Brunner, H.; Leyrer, H., Asymmetrische Katalysen: XXXV. Neue optisch aktive, einzähnige Phosphine aus  $\alpha$ -Pinen und Mannit. Rhodium-Komplexe und enantioselektive Katalysen. *J. Organomet. Chem.* **1987**, *334*, 369-376.
47. Apple, D. C.; Brady, K. A.; Chance, J. M.; Heard, N. E.; Nile, T. A., Iridium complexes as hydrosilylation catalysts. *J. Mol. Catal.* **1985**, *29*, 55-64.
48. Lavastre, O.; Morken, J. P., 500  $\mu$ m diameter beads as single reactors to screen organometallic catalysts: discovery of a new supported catalyst for the hydrosilylation of ketones. *New J. Chem.* **2002**, *26*, 745-749.
49. Boebel, T. A.; Hartwig, J. F., Silyl-Directed, Iridium-Catalyzed ortho-Borylation of Arenes. A One-Pot ortho-Borylation of Phenols, Arylamines, and Alkylarenes. *J. Am. Chem. Soc.* **2008**, *130*, 7534-7535.
50. Simmons, E. M.; Hartwig, J. F., Iridium-Catalyzed Arene Ortho-Silylation by Formal Hydroxyl-Directed C-H Activation. *J. Am. Chem. Soc.* **2010**, *132*, 17092-17095.
51. Ma, X.; Kucera, R.; Goethe, O. F.; Murphy, S. K.; Herzon, S. B., Directed C-H Bond Oxidation of (+)-Pleuromutilin. *J. Org. Chem.* **2018**, *83*, 6843-6892.
52. Kroke, E.; Li, Y. L.; Konetschny, C.; Lecomte, E.; Fasel, C.; Riedel, R., Silazane derived ceramics and related materials. *Materials Science & Engineering R-Reports* **2000**, *26*, 97-199.
53. Ionescu, E.; Kleebe, H. J.; Riedel, R., Silicon-containing polymer-derived ceramic nanocomposites (PDC-NCs): preparative approaches and properties. *Chem. Soc. Rev.* **2012**, *41*, 5032-5052.
54. Yilgor, I.; McGrath, J. E., Polysiloxane containing copolymers - a survey of recent developments. *Adv. Polym. Sci.* **1988**, *86*, 1-86.
55. Mark, J. E., Some interesting things about polysiloxanes. *Acc. Chem. Res.* **2004**, *37*, 946-953.
56. Bernoud, E.; Oulié, P.; Guillot, R.; Mellah, M.; Hannedouche, J., Well-Defined Four-Coordinate Iron(II) Complexes For Intramolecular Hydroamination of Primary Aliphatic Alkenylamines. *Angew. Chem. Int. Ed.* **2014**, *53*, 4930-4934.
57. King, A. K.; Buchard, A.; Mahon, M. F.; Webster, R. L., Facile, Catalytic Dehydrocoupling of Phosphines Using  $\beta$ -Diketimate Iron(II) Complexes. *Chem. Eur. J.* **2015**, *21*, 15960-15963.
58. Espinal-Viguri, M.; King, A. K.; Lowe, J. P.; Mahon, M. F.; Webster, R. L., Hydrophosphination of Unactivated Alkenes and Alkynes Using Iron(II): Catalysis and Mechanistic Insight. *ACS Catal.* **2016**, *6*, 7892-7897.

59. Espinal-Viguri, M.; Woof, C. R.; Webster, R. L., Iron-Catalyzed Hydroboration: Unlocking Reactivity through Ligand Modulation. *Chem. Eur. J.* **2016**, *22*, 11605-11608.
60. Coles, N. T.; Mahon, M. F.; Webster, R. L., Phosphine- and Amine-Borane Dehydrocoupling Using a Three-Coordinate Iron(II)  $\beta$ -Diketiminato Precatalyst. *Organometallics* **2017**, *36*, 2262-2268.
61. King, A. K.; Gallagher, K. J.; Mahon, M. F.; Webster, R. L., Markovnikov versus anti-Markovnikov Hydrophosphination: Divergent Reactivity Using an Iron(II)  $\beta$ -Diketiminato Pre-Catalyst. *Chem. Eur. J.* **2017**, *23*, 9039-9043.
62. Espinal-Viguri, M.; Neale, S. E.; Coles, N. T.; Macgregor, S. A.; Webster, R. L., Room Temperature Iron-Catalyzed Transfer Hydrogenation and Regioselective Deuteration of Carbon–Carbon Double Bonds. *J. Am. Chem. Soc.* **2019**, *141*, 572-582.
63. Reuter, M. B.; Cibazar, M. P.; Hammerton, J.; Waterman, R. Y., - Photoactivated silicon–oxygen and silicon–nitrogen heterodehydrocoupling with a commercially available iron compound. *Dalton Trans.* **2020**, *49*, 2972-2978.
64. See supporting information.
65. Vela, J.; Vaddadi, S.; Cundari, T. R.; Smith, J. M.; Gregory, E. A.; Lachicotte, R. J.; Flaschenriem, C. J.; Holland, P. L., Reversible Beta-Hydrogen Elimination of Three-Coordinate Iron(II) Alkyl Complexes: Mechanistic and Thermodynamic Studies. *Organometallics* **2004**, *23*, 5226-5239.
66. Dugan, T. R.; Bill, E.; MacLeod, K. C.; Brennessel, W. W.; Holland, P. L., Synthesis, Spectroscopy, and Hydrogen/Deuterium Exchange in High-Spin Iron(II) Hydride Complexes. *Inorg. Chem.* **2014**, *53*, 2370-2380.
67. The analogous iron-deuteride dimer is also observed by  $^2\text{H}$  NMR during KIE studies.
68. Yu, Y.; Brennessel, W. W.; Holland, P. L., Borane B–C Bond Cleavage by a Low-Coordinate Iron Hydride Complex and N–N Bond Cleavage by the Hydridoborate Product. *Organometallics* **2007**, *26*, 3217-3226.
69. See supporting information.
70. Evans, R.; Deng, Z.; Rogerson, A. K.; McLachlan, A. S.; Richards, J. J.; Nilsson, M.; Morris, G. A., Quantitative Interpretation of Diffusion-Ordered NMR Spectra: Can We Rationalize Small Molecule Diffusion Coefficients? *Angew. Chem. Int. Ed.* **2013**, *52*, 3199-3202.
71. Ekkert, O.; White, A. J. P.; Toms, H.; Crimmin, M. R. Y., - Addition of aluminium, zinc and magnesium hydrides to rhodium(iii). *Chem. Sci.*, **2015**, *6*, 5617-5622.